Black Cherry

Prunus serotina





Mature

General	Attn	ihutaa
General	Au	inuies

Type Deciduous Tree Height 70 - 100 Feet **Spread** 35 - 50 Feet **Form** Rounded **Utility Lines** Incompatible

Growth Rate Fast Life Expectancy Medium **USDA** Zone 3 - 9

Root Pattern Deep Lateral



Native Range

Plant Community

Sun Full Sun, Partial Sun Orientation North, East

Soil Texture Sandy Loam to Clay

Topography Upland

Forest, Open Cliff, Savannah, Forest Edge, Old Field **Plant Community**

Succession Pioneer

Origin MSA Zone 1, MSA Zone 2,

> MSA Zone 3, MSA Zone 4, MSA Zone 5, MSA Zone 6, Minnesota, North America

Flowers/Foliage/Fruits

Flower Color White Flower Season Spring Fruit Drupe **Fruit Color** Black **Fruit Season** Summer **Summer Texture** Medium **Winter Texture** Fine **Spring Foliage** Green **Summer Foliage** Green **Fall Foliage** Yellow

Winter Foliage Not Applicable

Soils

Salt Spray Tolerance Moderate **Soil Salt Tolerance Tolerant Compaction Tolerance** Intolerant Water Table 24 Inches

Drainage Excessive, Moderate

Flood Tolerance

Drought Tolerance Tolerant Moisture Regime Moist, Wet 6.0 through 7.5 pН

Windbreak Group

Black Cherry

Prunus serotina



P	lan	tıng	Ease

Spring Bareroot Fall Bareroot

Spring Container
Fall Container

Spring Seed Fall Seed

Maintenance

Formal Moderate
Informal Low
2,4-D Tolerance Sensitive
Dicamba Tolerance Sensitive
Picloram Tolerance Sensitive

Clopyralid Tolerance Artificial Light Sulfur Dioxide Ozone Tolerance

Hydrogen Flouride Moderate

Nitrogen Oxide

Pests/Problems

Allergens both sexes 5-7 depending on

species depending on variety

Invasiveness Invasive

Toxicity Has toxic Roots, Stems, Leaves,

Easy

Easy

Easy

Flowers, Fruit

Cold InjuryNot SusceptibleStorm DamageSusceptible

Biological Control Mechanical Control Chemical Control

Miscellaneous

Wildlife Rating High
Insect Concerns Minor
Disease Concern Minor

Wildlife Concerns

Comments

A native tree valued for its lumber. Bark smooth and brown becoming red-gray, shaggy, and peeling on older trees. Bark, leaves, and fruit kernels contain cyanide.



Bark



Dormant











Flower



Leaf